



## BANWEED MCPA

REG. NO. L0466 / N-AR 0121  
W 130201

ACT No. 36 OF 1947

A SOLUBLE CONCENTRATE HERBICIDE FOR THE CONTROL OF ANNUAL BROADLEAF WEEDS, AS LISTED, IN LAWNS AND TURF	'N OPLOSBARE KONSENTRAAT- ONKRUIDDODER VIR DIE BEHEER VAN EENJARIGE BREëBLAARONKRUID, SOOS AANGETOON, IN GRASPERKE EN GRASVELDE	
HRAC HERBICIDE GROUP CODE	0	HRAC ONKRUIDDODER GROEP KODE

### Active Ingredient / Aktiewe Bestanddeel

MCPA (Phenoxyacetic acid) .....400 g/l .....MCPA (Fenoksieasynsuur)  
(as Potassium salt) (as Kaliumsout)

### NETT VOLUME / NETTO VOLUME

Registered by: / Geregistreer deur:  
Agro-Serve (Pty) Ltd t/a Efekto™ Reg. No. 1973/000868/07  
Agro-Serve (Edms) Bpk h/a Efekto  
Distributed by: / Versprei deur:  
Efekto Care (Pty) Ltd Reg. No.2009/006357/07 Efekto Care (Edms) Bpk.  
PO Box / Posbus 652147, BENMORE. 2010  
Efekto and the EFEKTO logo are trademarks of Agro-Serve (Pty) Ltd  
Efekto en die EFEKTO kenteken is handelsmerke van Agro-Serve (Edms) Bpk  
© Agro-Serve (Pty) Ltd. © Agro-Serve (Edms) Bpk

All rights reserved / Alle regte voorbehou  
CUSTOMER CARE: Tel.0861 EFEKTO (333586) office hours / kantoor ure  
[info@efekto.co.za](mailto:info@efekto.co.za) / [www.efekto.co.za](http://www.efekto.co.za)

EMERGENCY NUMBER / NOODNOMMER  
083 1233911

**BEFORE USING THIS PRODUCT PLEASE READ the LABEL CAREFULLY**

BATCH NUMBER / LOTNOMMER

DATE MANUFACTURED / DATUM VERVAARDIG

UN NUMBER: 3082



**HARMFUL / SKADELIK**

## **WAARSKUWINGS**

Hanteer versigtig.

Giftig as ingesluk of ingeasem word of met die vel in aanraking kom.

Bêre in 'n koel plek.

Bêre weg van voedsel, voer, saad en ander landboumiddels en buite bereik van kinders, oningeligte persone en diere.

## **VERHOED SPUITSTOFWEGDRYING NA GEVOELIGE PLANTE, BYVOORBEELD ALLE BREËBLAARGEWASSE ASOOK ALLE BOME, STRUIKE EN BLOMPLANTE.**

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie, aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale klimaats- en bergingstoestande; kwaliteit van verdunningswater; verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die onkruid teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

## **VOORSORGMAATREËLS**

Vermyn inaseming van sproeireën en spuitdampe.

Was besoedelde klere daagliks.

Was onmiddellik met seep en water indien per ongeluk met die vel in aanraking kom.

Moenie tydens of na vermenging en toediening eet, drink of rook alvorens hande en gesig gewas en ander klere aangetrek is nie.

Verhoed spuitstofwegdrywing na ander gewasse, weiding, riviere, damme en enige gebied wat nie behandel word nie.

Om beskadiging van gewenste plante te voorkom, moet spuittoerusting wat vir toediening gebruik is, nie weer gebruik word vir toediening van ander middels op sulke plante nie.

Gooi was- en spoelwater weg waar dit nie gewasse, weiding, riviere en damme sal besoedel nie.

Spoel leë houer driekeer uit met 'n volume water gelykstaande aan 'n minimum van 10 % van die houer. Gooi die spoelwater by die inhoud van die spuittenk.

Vernietig leë houer. Moet nooit vir enige ander doel gebruik nie.

Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.

## SOMMIGE ONKRUIDSOORTE WAT NORMAALWEG BEHEER WORD

<i>Arctotis venusta</i>	gousblom
<i>Bidens pilosa</i>	knapsekêrel
<i>Chenopodium album</i>	withondebossie
<i>Cosmos bipinnatus</i>	kosmos
<i>Galinsoga parviflora</i>	knopkruid
<i>Raphanus raphanistrum</i>	ramnas
<i>Striga asiatica</i>	rooiblom
<i>Tagetes minuta</i>	kakiebos
<i>Tribulus terrestris</i>	dubbeltjie
<i>Xanthium spinosum</i>	boetebossie

## GEBRUIKSAANWYSINGS AANGEDUI

## GEBRUIK SLEGS SOOS

Dien toe na opkom van breëblaaronkruid wat in gevestigde grasperke en grasvelde groei.

Dien toe as 'n egalige fyn bespuiting.

Kalibreer spuittoerusting noukeurig voor toediening.

## LET WEL:

**EFEKTO BANWEED MCPA** mag 'n verkleuring van die grasperk veroorsaak.

Dit sal egter tydelik van aard wees.

Die toediening van 'n stikstofhoudende kunsmis 2 - 3 weke voor toediening van die onkruidodder word aanbeveel.

## VERMENGING EN TOEDIENING

Verdun 50 ml **EFEKTO BANWEED MCPA** in 5 l water en dien egalig toe oor 100 m<sup>2</sup>.

Herhaal bespuiting na 14 dae indien nodig.

Gebruik slegs op onkruid wat aktief groei in grond met voldoende vog.

Gebruik slegs op die skoonveld in die geval van gholfbane.

## BELANGRIK

Toediening deur middel van gieters of toerusting wat aan tuinslange gekoppel word, word nie aanbeveel nie.

Rugsakspuite en drukspuitpompe is egter vir die doel geskik.

## **WARNINGS**

Handle with care.

Poisonous when absorbed through the skin, swallowed or inhaled.

Store in a cool place.

Store away from food, feed, seed, other agricultural remedies and out of reach of children, uninformed persons and animals.

**AVOID SPRAY DRIFT ONTO SUSCEPTIBLE PLANTS E.G. ALL BROAD LEAF CROPS AS WELL AS ALL TREES, SHRUBS AND FLOWERING PLANTS.**

**Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label and occurrence of resistance of the weed against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.**

## **PRECAUTIONS**

Do not inhale spray mist and fumes.

Wash contaminated clothing daily.

Wash with soap and water immediately after accidental skin contact.

Do not eat, drink or smoke whilst mixing and applying or before washing hands and face and change of clothing.

Avoid spray drift onto other crops, grazing, rivers, dams and any area not under treatment.

To avoid injury to desirable plants, spray equipment used for application should not be re-used to apply other materials to such plants.

Dispose of wash water where it will not contaminate crops, grazing, rivers and dams.

Rinse empty container three times with a volume of water equal to a minimum of 10 % of that of the container. Add the rinse water to the contents of the spray tank.

Destroy empty container. Never use for any other purpose.

Prevent contamination of food, feeds, drinking water and eating utensils.

## SOME WEED SPP. NORMALLY CONTROLLED

<i>Arctotis venusta</i>	Free State daisy
<i>Bidens pilosa</i>	blackjack
<i>Chenopodium album</i>	white goosefoot
<i>Cosmos bipinnatus</i>	cosmos
<i>Galinsoga parviflora</i>	small flowered quick weed
<i>Raphanus raphanistrum</i>	wild radish
<i>Striga asiatica</i>	witchweed
<i>Tagetes minuta</i>	khaki weed
<i>Tribulus terrestris</i>	common dubbeltjie
<i>Xanthium spinosum</i>	spiny cockle bur

## DIRECTIONS FOR USE DIRECTED

## USE ONLY AS

Apply after emergence of broadleaf weeds growing in an established lawn or turf.  
Apply as a fine, even spray.  
Calibrate spray equipment carefully prior to application.

## NOTE

**EFEKTO BANWEED MCPA** may cause a discoloration of the grass.

This will be of a temporary nature only.

The application of a nitrogenous fertiliser 2 - 3 weeks prior to treating with the herbicide is recommended.

## MIXING AND APPLICATION

Dilute 50 ml **EFEKTO BANWEED MCPA** in 5 l water and apply as an even spray onto 100 m<sup>2</sup>.

Repeat application after 14 days if necessary.

Use only on weeds growing actively in a soil with sufficient moisture.

Use only in fairways in the case of golf courses.

## IMPORTANT

Application by means of a watering can or equipment attached to garden hoses is not recommended.

Knapsack sprayers and pressurised sprayers are however suitable equipment to use.