



FUNGINEX

REG. NO. L 2469 / N-AR 0131
W 130281

AN EMULSIFIABLE CONCENTRATE
SYSTEMIC FUNGICIDE FOR THE
CONTROL OF POWDERY MILDEW,
BLACK SPOT AND RUST ON VARIOUS
ORNAMENTALS AND POWDERY
MILDEW AND RUST ON SEVERAL
EDIBLE CROPS

ACT NO 36 of 1947

'N EMULGEERBARE KONSENTRAAT
SISTEMIESE SWAMDODER VIR DIE
BEHEER VAN POEIERAGTIGE
SKIMMEL, SWARTVLEK EN
BRUINROES OP VERSKILLENDE
SIERPLANTE EN POEIERAGTIGE
SKIMMEL EN BRUINROES OP
VERSKEIE EETBARE GEWASSE

FRAC FUNGICIDE GROUP CODE

3

FRAC SWAMDODER GROEP KODE

Active Ingredient / Aktiewe Bestanddeel

Triforine.....190 g/l.....Triforien

NET VOLUME / NETTO VOLUME

Registered by: / Geregistreer deur:

Agro-Serve (Pty) Ltd t/a Efekto™ Reg. No. 1973/000868/07 Agro-Serve (Edms) Bpk h/a Efekto

Distributed by: / Versprei deur:

Efekto Care (Pty) Ltd Reg. No.2009/006357/07 Efekto Care (Edms) Bpk.
PO Box / Posbus 652147, BENMORE. 2010

Efekto and the EFEKTO logo are trademarks of Agro-Serve (Pty) Ltd
Efekto en die EFEKTO kenteken is handelsmerke van Agro-Serve (Edms) Bpk

© Agro-Serve (Pty) Ltd. © Agro-Serve (Edms) Bpk

All rights reserved / Alle regte voorbehou

CUSTOMER CARE: Tel.0861 EFEKTO (333586) office hours / kantoor ure

info@efekto.co.za / www.efekto.co.za

©Funginex is the registered trademark of Sumitomo (Pty) Ltd.

EMERGENCY NUMBER / NOODNOMMER

083 1233911

BEFORE USING THIS PRODUCT PLEASE READ the LABEL CAREFULLY

BATCH NUMBER / LOTNOMMER

DATE MANUFACTURED / DATUM VERVAARDIG

UN NUMBER: Not regulated



HARMFUL / SKADELIK



WARNINGS

THE FOLLOWING NUMBER OF DAYS MUST BE ALLOWED BETWEEN LAST APPLICATION AND HARVEST OR FEEDING OF CROP

CUCURBITS	1 day
BEANS	3 days
PEAS	4 days

Keep out of reach of children, uninformed persons and animals.
Store away from food and feed.
FLAMMABLE - Keep away from fire.

PRECAUTIONS

Do not breathe spray mist and avoid skin contact.
Do not eat, drink or smoke whilst applying or before washing hands and face.
Avoid contamination of food, eating utensils, drinking water, other edible crops and fishponds.
Destroy empty container and do not re-use for any other purpose.

SYMPTOMS OF HUMAN POISONING

Headache, nausea, vomiting, anxiety, muscle twitching beginning in eye lids and tongue.

FIRST AID TREATMENT

Eye contact:

Flush eyes immediately with fresh water for at least 15 minutes while holding the eyelids open.
Remove contact lenses if worn. Call a physician.

Skin contact:

If on skin, remove contaminated clothing, wash skin with plenty of soap and water. Wash contaminated clothing before reuse.

Ingestion:

If swallowed, give water or milk to drink. DO NOT make person vomit unless directed to do so by medical personnel. If medical advice cannot be obtained, then immediately take the person and product container, with label, to the nearest medical emergency treatment center or hospital.

Inhalation:

If respiratory discomfort or irritation occurs, move the person to fresh air. See a doctor if discomfort or irritation continues.

NOTE TO PHYSICIAN

This product is of very low acute toxicity and poisoning is unlikely to occur. Treatment should be symptomatic.

DIRECTIONS FOR USE

USE ONLY AS DIRECTED

Apply as a full cover spray to the foliage either as preventative treatment or when the first symptoms are observed.

Repeat at weekly intervals to keep plants free of disease.

The shorter spray intervals should be used when conditions are especially favourable for infection.

Do not spray into open flowers.

CROP	DOSAGE / 10 ℓ WATER	REMARKS
ROSES 1. In the open a) Powdery mildew b) Black spot, Rust 2. In shade All diseases mentioned	10 mℓ 12,5 mℓ 20 mℓ	Apply every 7 - 10 days or when necessary. Wet plants thoroughly
FLOWERS & ORNAMENTALS Powdery mildew	15 mℓ	Repeat when necessary.
CARNATIONS Rust	12,5 mℓ	Repeat weekly when necessary.
CUCURBITS Powdery mildew	12,5 mℓ	Repeat every 7 - 10 days or when necessary. Wet plants thoroughly.
BEANS AND GREEN BEANS Rust	15 mℓ	Commence spraying at the first signs of infection. Repeat every 7 - 10 days if necessary. Wet plants thoroughly.
PEAS Powdery mildew	15 mℓ	Repeat spray every 7 - 10 days if necessary. Under conditions of severe infection use the shorter spray interval.

WAARSKUWINGS

LAAT DIE VOLGENDE AANTAL DAE VERLOOP TUSSEN DIE LAASTE BESPUITING EN OES:

PAMPOENGWASSE	1 dag
BOONTJIES	3 dae
ERTJIES	4 dae

Hou buite bereik van kinders, oningeligte persone en diere.

Bêre weg van kos en veevoer.

VLAMBAAR - Hou weg van vuur.

VOORSORGMAATREËLS

Voorkom inaseming van spuitmis en velkontak.

Moenie eet, drink of rook tydens toediening of voordat gesig en hande gewas is nie.

Voorkom besoedeling van voedsel, eetgerei, drinkwater, ander eetbare gewasse en visdammetjies.

Vernietig leë houers en moet vir geen ander doel gebruik nie.

SIMPTOME VAN MENSLIKE VERGIFTIGING

Hoofpyn,naarheid,braking, angstigheid, spierbewegings in die ooglede en tong,

NOODHULP BEHANDELING

Oogkontak:

Spoel oë onmiddelik met skoon water vir ten minste 15 minute terwyl die ooglede oop gehou word.

Verwyder kontaklense. Raadpleeg 'n geneesheer.

Velkontak:

Wanneer in kontak gemaak is met die vel, verwyder gekontameneerde klere, was vel met skoon water en seep. Was gekontameneerde kleding voor hergebruik.

Inname:

Wanneer ingesluk, gee water of melk om te drink. MOENIE persoon laat braak tensy voorgestel deur mediese personeel. Indien mediese persoon nie beskikbaar is nie, neem pasient met die houer en die etiket na die naaste geneesheer of mediese ongevalle.

Inaseming:

Indien respiratoriese ongemak of irritasie voorkom, verwyder persoon na vars lug. Raadpleeg 'n geneesheer indien ongemak of irritasie voortduur.

NOTA AAN GENEESHEER

Hierdie produk het 'n baie lae akute toksisiteit en vergiftiging is onwaarskynlik. Behandel simptome.

GEBRUIKSAANWYSINGS

GEBRUIK SLEGS SOOS AANGEDUI

Dien toe as 'n volle dekbespuiting op die blare of as 'n voorkomende behandeling of wanneer eerste simptome opgemerk word.

Herhaal met weeklikse tussenposes om plante vry van siekte te hou.

Die korter periodes tussen toedienings behoort gebruik te word wanneer die omstandighede besonder gunstig vir besmetting is.

Moet nie in oop blomme spuit nie.

GEWAS	DOSIS / 10 ℓ WATER	OPMERKINGS
ROSE 1. In opelug a) Poeieragtige skimmel b) Swartvlek, Bruinroes 2. Onder skadudoek Alle siektes soos vermeld	10 mℓ 12,5 mℓ 20 mℓ	Herhaal elke 7 - 10 dae of wanneer nodig. Benat plante deeglik
BLOMME & SIER-PLANTE Poeieragtige skimmel	15 mℓ	Herhaal wanneer nodig.
ANGELIERE Bruinroes	12,5 mℓ	Herhaal weekliks indien nodig.
PAMPOENGWASSE Poeieragtige skimmel	12,5 mℓ	Herhaal elke 7 - 10 dae of wanneer nodig. Benat die plant deeglik.
BONE EN GROENBONE Bruinroes	15 mℓ	Begin spuit sodra eerste tekens van besmetting waargeneem word. Herhaal elke 7 - 10 dae indien nodig. Benat die plant deeglik.
ERTJIES Poeieragtige skimmel	15 mℓ	Herhaal bespuiting elke 7 - 10 dae indien nodig. Gebruik die korter spuit tussenpose gedurende

		strawwe besmettings.
--	--	----------------------