

KAMIKAZE (KLEIN VERPAKKING)



DRAFT LABEL / KONSEP ETIKET

KAMIKAZE

REG. NO. L 4216 / N-AR 0106
W130194

ACT NO 36 OF 1947

A BAIT FOR THE CONTROL OF THE
HARVESTER TERMITE

'N LOKAAS VIR DIE BEHEER VAN DIE
GRASDRAERTERMIET

IRAC INSECTICIDE GROUP CODE 1A INSEKDODER GROEP KODE

Active Ingredient / Aktiewe Bestanddeel

Carbaryl (Carbamate).....16,0 g/kg.....Karbaril (Karbamaat)

NET MASS / NETTO MASSA

Registered and Distributed by / Geregistreer en Versprei deur:
Agro-Serve (Pty) Ltd trading as EFEKTO™ Reg. No. 1973/000868/07 Private Bag X 215, Bryanston, 2021
EFEKTO and the EFEKTO logo are trademarks of Agro-Serve (Pty) Ltd.
© Agro-Serve (Pty) Ltd. All rights reserved.

CUSTOMER CARE / KLIËNTEDIENS: Tel: 0861 EFEKTO (33586) Office Hours / Kantoor ure
info@efekto.co.za / www.efekto.co.za
EMERGENCY NUMBER/NOODNOMMER 0831233911

BEFORE USING THIS PRODUCT PLEASE READ the LABEL CAREFULLY

BATCH NUMBER / LOTNOMMER

DATE MANUFACTURED / DATUM VERVAARDIG

UN NUMBER: 2757



WAARSKUWINGS

ONTTREK LEWENDEHAWE TYDENS EN VIR 'N TYDPERK VAN DRIE DAE NA TOEDIENING.

Giftig vir vis en pluimvee.

Berg in 'n koel plek weg van voedsel en voer.

Hou buite bereik van kinders, oningeligte persone en diere.

VOORSORGMAATREËLS

Moet nie eet, drink of rook gedurende toediening of voordat hande en gesig gewas is nie.

Vermý besoedeling van visdammetjies, riviere en damme.

Was met seep en water na hantering.

Vernietig leë houer en moet vir geen ander doel gebruik nie.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale grond-, klimaats- en bergingstoestande; kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket is nie en die voorkoms van weerstand van die plaag teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse plantegroei, die omgewing of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

GEBRUIKSAANWYSINGS

GEBRUIK SLEGS SOOS AANGEDUI

Strooi die aas eweredig oor die besmette gebied en rondom aktiefwerkende nelsingange. Die aangewese tyd om die aas uit te strooi is gedurende Mei, Junie, Julie en Augustus. Gewoonlik is die termiete op hulle bedrywigste vanaf 11 uur soggens tot sonder. Strooi die aas op die dae waarop intense termietbedrywigheid voorkom. Een toediening is normaalweg voldoende om termietbeheer te verkry. Strooi slegs soveel aas as wat die termiete op dag van behandeling nes toe kan dra. Die resultaat moet na een maand bepaal word en die plekke waar termietbedrywigheid nog voorkom, moet weer met lokaas behandel word.

VERMY NAT TOESTANDE, WANT REËN VERNIETIG AAS. AAS WERK AFWEREND WANNEER BESOEDEL MET SKERPRUIKENDE STOWWE SOOS VISMEEL, VETTE, OLIES, PETROL EN SELFS DIE REUK VAN 'N MENS. DIT IS DAAROM GERADE OM RUBBER OF PVC HANDSKOENE TE DRA WANNEER U DIE AAS HANTEER.

WARNINGS

WITHDRAW LIVESTOCK FROM TREATED AREAS DURING AND FOR A PERIOD OF THREE DAYS AFTER TREATMENT.

Toxic to fish and poultry.

Store in a cool place away from food and feed.

Keep out of reach of children, uninformed persons and animals.

PRECAUTIONS

Do not eat, drink or smoke whilst applying, before washing hands and face.

Avoid contamination of fish ponds, rivers and dams.

Wash with soap and water after handling.

Destroy empty container and do not use for any other purposes.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty

DIRECTIONS FOR USE

USE ONLY AS DIRECTED

Scatter bait evenly over infested area and around actively working nest entrances. The best period in which to scatter bait is during May, June, July and August. Usually the termites are active from 11:00 until sundown. Scatter the bait on a day when termites are intensely foraging. One application will normally suffice to control the termites. Only scatter as much bait as the termites can carry to their nests on the day of treatment. The result of the application should be assessed after one month and if activity is still occurring, repeat application.

AVOID WET CONDITIONS, BECAUSE RAIN DESTROYS THE BAIT. BAIT WILL REPEL TERMITES IF CONTAMINATED WITH MINUTE QUANTITIES OF SHARP-SMELLING SUBSTANCES SUCH AS FISHMEAL, FATS, OILS, PETROL AND EVEN THE SMELL OF HUMAN BEINGS. IT IS THEREFORE ADVISABLE TO WEAR RUBBER OR PVC GLOVES WHEN HANDLING THE BAIT.