



# KARBADUST

REG. NO. L 0493 & G 0196 / N-AR 0148  
W130190

ACT NO 36 OF 1947

INSECTICIDE DUSTING POWDER FOR  
HOME GARDEN AND ANIMAL USE. A  
RESIDUAL CONTACT AND STOMACH  
POISON FOR THE CONTROL OF PESTS  
AS LISTED

INSEKDODER-STUIFPOEIER VIR  
HUISTUIN- EN DIERE GEBRUIK. 'N  
KONTAK- EN MAAGGIF MET NAWERKING  
VIR DIE BEHEER VAN PLAE SOOS  
AANGEDUI

STORE IN A COOL DRY PLACE / BERG IN 'n KOEL DROë PLEK

**IRAC INSECTICIDE GROUP CODE 1 A INSEKDODER GROEP KODE**  
**ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL**

Carbaryl (Carbamate)..... 50 g/kg.....Karbaril (Karbamaat)

## NET MASS / NETTO MASSA

Registered by: / Geregistreer deur:

Agro-Serve (Pty) Ltd t/a Efekto™ Reg. No. 1973/000868/07 Agro-Serve (Edms) Bpk h/a Efekto

Distributed by: / Versprei deur:

Efekto Care (Pty) Ltd Reg. No.2009/006357/07 Efekto Care (Edms) Bpk.

PO Box / Posbus 652147, BENMORE. 2010

Efekto and the EFEKTO logo are trademarks of Agro-Serve (Pty) Ltd

Efekto en die EFEKTO kenteken is handelsmerke van Agro-Serve (Edms) Bpk

© Agro-Serve (Pty) Ltd. © Agro-Serve (Edms) Bpk

All rights reserved / Alle regte voorbehou

CUSTOMER CARE: Tel.0861 EFEKTO (333586) office hours / kantoor ure

[info@efekto.co.za](mailto:info@efekto.co.za) / [www.efekto.co.za](http://www.efekto.co.za)

**EMERGENCY NUMBER / NOODNOMMER**

**083 1233911**

**BEFORE USING THIS PRODUCT PLEASE READ the LABEL CAREFULLY**

**BATCH NUMBER / LOTNOMMER**

**DATE MANUFACTURED / DATUM VERVAARDIG**

**UN NUMBER: 2757**



**CAUTION / VERSIGTIG**



## **WAARSKUWINGS**

LAAT 14 DAE VERLOOP TUSSEN LAASTE TOEDIENING EN OES OF BEWEIDING EN 7 DAE TUSSEN LAASTE AANWENDING EN SLAG VAN HOENDERS VIR MENSLIKE GEBRUIK.

Giftig deur inname en inaseming.

Kan oë irriteer.

Giftig vir visse, bye en wild.

Berg weg van voedsel en voer.

Hou buite bereik van kinders, oningeligte persone en diere.

Hou houer dig toe.

## **VOORSORGMAATREËLS**

Vermy langdurige kontak met, of inaseming van poeier.

Was deeglik met seep en water na gebruik of toevallige velkontak.

Moenie eet, drink of rook tydens toediening nie.

Voorkom dat poeier oorwaai na ander gewasse of dele wat nie onder behandeling is nie.

Vernietig leë houer en moet dit vir geen ander doel gebruik nie.

Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.

## **GEBRUIKSAANWYSINGS**

## **GEBRUIK SLEGS SOOS AANGEDUI**

### **LUSERNRUSPES, GRASPERKRUSPES, KOMMANDOWURMS, STINKSPRINKANE EN KRIEKE OP GRASPERKE**

Karbadust-stuifpoeier moet teen 20 g / 10 m<sup>2</sup> toegedien word in gebiede waar hierdie plaehul verskyning maak.

Waar grasperkruspes voorkom, moet die grasperk na bestuiwing liggies natgespuit word.

Waar krieke voorkom moet daar binne-in die gate bestuif word.

Een toediening is gewoonlik voldoende.

## **SKILDSTINKBESIES, LENTEKEWERS EN ASTYLUSKEWERS**

Bestuif plante deeglik sodra hierdie plaë hulle verskyning maak.

## **ROOIMYTE EN TAMPANS BY HOENDERS EN KOUVOËLS.**

Bestuif alle hokke deeglik en skenk veral aandag aan die vloer, nuwe skropgoed, stellasies, neste, nuwe nesstrooi en barsies in die houtwerk en mure.

## **LUISE**

Sprinkel die poeier onder albei vlerke en onder die stert.

## **BOSLUISE EN VLOOIE OP HONDE EN KATTE**

Sprinkel die poeier liggies op die hare en vryf deeglik in. By katte moet die poeier weer deeglik uitgeborsel word. Behandel weekliks of wanneer dit nodig is.

## **WARNINGS**

ALLOW 14 DAYS BETWEEN LAST APPLICATION AND HARVEST OR GRAZING AND 7 DAYS BETWEEN LAST APPLICATION AND SLAUGHTER OF FOWLS FOR HUMAN CONSUMPTION.

Poisonous by swallowing and inhalation.

May irritate eyes.

Toxic to fish, bees and wildlife.

Store away from food and feed.

Keep out of reach of children, uninformed persons and animals.

Keep container well closed.

## **PRECAUTIONS**

Avoid prolonged contact with, or inhalation, of dust.

Wash thoroughly with soap and water after use or accidental skin contact.

Do not eat, drink or smoke whilst applying.

Prevent drift onto other crops or areas not under treatment.

Destroy empty container and do not use for any other purpose.

Prevent contamination of food, feed, drinking water and eating utensils.

## **DIRECTIONS FOR USE**

## **USE ONLY AS DIRECTED**

### **LUCERNE CATERPILLARS, LAWN CATERPILLARS, ARMY WORMS, ELEGANT GRASS-HOPPERS AND CRICKETS ON LAWNS**

Karbadust dusting powder must be applied at the rate of 20 g / 10 m<sup>2</sup> on areas infested by these pests.

Where lawn caterpillars occur the lawn must be lightly irrigated after dusting.

Where crickets occur, dust inside the holes.

One application should suffice.

### **SHIELD BUGS, CHAFER BEETLES AND ASTYLUS BEETLES**

Dust plants thoroughly when these pests are noticed.

### **RED MITES AND TAMPANS ON POULTRY AND CAGE BIRDS**

Dust all poultry houses thoroughly and pay particular attention to the floor, new litter, perches, nests and nest hay as well as cracks in woodwork and walls.

### **LICE**

Dust under each wing and the vent.

### **TICKS AND FLEAS ON DOGS AND CATS**

Sprinkle the powder lightly on the hair and rub in well.  
In the case of cats the powder must be brushed out afterwards.  
Apply treatment weekly or as often as is necessary.