



ROSECARE 3

REG. NO. L 6509 N-AR 0142
W130279

ACT NO 36 OF 1947

A CONTACT INSECTICIDE PLUS A SYSTEMIC FUNGICIDE FORMULATED AS A MICRO EMULSION FOR THE CONTROL OF INSECTS AND FUNGUS DISEASES AS INDICATED ON ROSES AND OTHER ORNAMENTALS

'N KONTAK INSEKDODER PLUS 'N SISTEMIESE SWAMDODER GEFORMULEER AS 'N MIKRO EMULSIE VIR DIE BEHEER VAN INSEKTE EN SWAMSIKTES SOOS AANGEDUI OP ROSE EN ANDER SIERPLANTE

FRAC FUNGICIDE GROUP CODE	3	FRAC SWAMDODER GROEP KODE
IRAC INSECTICIDE GROUP CODE	3 A	IRAC INSEKDODER GROEP KODE

Active Ingredients / Aktiewe Bestanddele

Bifenthrin(synthetic pyrethroid)..... 2.0 g/l Bifentrin(sintetiese piretroied)
Myclobutanil(triazole) 7.5 g/l Miklobutaniel(triasool)

NET VOLUME / NETTO VOLUME

Registered by / Geregistreer deur:
Agro-Serve (Pty) Ltd trading as EFEKTO™ Reg. No. 1973/000868/07
Distributed by; / Versprei deur:
Efekto Care (Pty) Ltd
Reg. No. 2009/00635/07
PO Box 652147, BENMORE, 2010
EFEKTO and the EFEKTO logo are trademarks of Agro-Serve (Pty) Ltd.
© Agro-Serve (Pty) Ltd. All rights reserved.
CUSTOMER CARE / KLIËNTEDIENS: Tel: 0861 EFEKTO (333586) OH/KU
info@efekto.co.za / www.efekto.co.za
EMERGENCY NUMBER/NOODNOMMER 0831233911
BEFORE USING THIS PRODUCT PLEASE READ the LABEL CAREFULLY
MANUFACTURED BY/ VERVAARDIG DEUR
THE SCOTTS COMPANY
BEFORE USING THIS PRODUCT PLEASE READ the LABEL CAREFULLY
BATCH NUMBER / LOTNOMMER

DATE MANUFACTURED / DATUM VERVAARDIG

UN NUMBER: 3082



VERSIGTIG / CAUTION



Beskerm rose en ander sierplante teen Poeieragtige skimmel, Swartvlek, Bruinroes, Plantluise. Blaaspootjies en Rooispinmyt

Rosecare 3 bevat 'n sistemiese swamdoder om genoemde siektes te voorkom en te beheer en 'n insekdoder om plantluise, blaaspootjies en roospinmyt effektief te beheer.

WAARSKUWINGS

- Moet nie toedien op eetbare gewasse nie.
- Skadelik indien deur die vel geabsorbeer, ingesluk of ingesem word.
- Giftig vir visse en ander waterorganismes
- Skadelik vir bye.
- Berg in 'n koel plek.
- Berg weg van voedsel en voer.
- Hou in oorspronklike houer, stewig toe, in 'n koel plek.
- Hou buite bereik van kinders, oningeligte persone en diere.

VOORSORGMAATREËLS

- Dra rubberhandskoene wanneer die konsentraat hanteer word.
- Vermy oog- en velkontak met die konsentraat.
- Voorkom inaseming van spuitnewel.
- Was met seep en water na gebruik.
- Maak houer heeltemal leeg en gooi op 'n veilige plek weg.
- Vermy besoedeling van voedsel, voer, drinkwater en eetgerei.
- Vermy wegdrywing van spuitmis na eetbare gewasse en visdammetjies.

GEBRUIKSAANWYSINGS

GEBRUIK SLEGS SOOS AANGEDUI

WANNEER :

Spuit elke 7 - 10 dae solank insekte en siektes opgemerk word. Blaaspootjies, Rooispinmyt en Poeieragtige skimmel sal slegs effektief beheer word as 'n spuitprogram van minstens 3 opeenvolgende bespuitings gedoen word. As swartvlek die vorige seisoen teenwoordig was moet bespuitings alreeds begin word wanneer die botsels begin oopmaak.

LET WEL: Indien daar alreeds Blaaspootjies, Rooispinmyt of Poeieragtige skimmel teenwoordig is moet minstens 3 bespuitings weekliks teen die hoër dosis gespuit word.

HOE :

- SKUD GOED VOOR GEBRUIK.
- Voeg 10 ml produk by elke 1 liter water.
- Indien Blaaspootjies, Rooispinmyt of Poeieragtige skimmel alreeds teenwoordig is voeg 15 ml produk by elke 1 liter water.
- Spuit deeglik nat.
- Was die spuit goed uit na gebruik.
- Moet nie oortollige mengsel bêre nie.

Protects roses and other ornamentals against Powdery mildew, Blackspot, Rust, Aphids, Thrips and Redspidermite.

Rosecare 3 contains a systemic fungicide to prevent **and** control diseases mentioned and an insecticide which controls aphids, thrips and redspidermites effectively.

WARNINGS

- Do not apply to edible crops.
- Harmful when absorbed through the skin, swallowed or inhaled.
- Toxic to fish and other aquatic organisms
- Harmful to bees.
- Store in a cool place.
- Store away from food and feed.
- Keep in original container, tightly closed, in a safe place.
- Keep out of reach of children, uninformed persons and animals.

PRECAUTIONS

- Wear rubber gloves when handling the concentrate.
- Avoid eye and skin contact with the concentrate.
- Do not breathe spray mist.
- Wash with soap and water after use.
- Empty container completely and dispose of safely.
- Avoid contamination of food, feed, drinking water and eating utensils.
- Avoid spray drift onto **edible** crops and fishponds.

WHEN :

Spray every 7 - 10 days as long as insects or diseases are noticed. Thrips, Redspidermite and Powdery mildew will only be controlled effectively if a spray programme of at least 3 successive sprays are done. When rose plants were infested with blackspot during the previous season, spraying should commence as soon as buds start to burst.

IMPORTANT: If Thrips, Redspidermite or Powdery mildew are already present at least 3

applications at the higher dosage rate should be done at weekly intervals.

HOW :

- SHAKE WELL BEFORE USE.
- Add 10 ml of product to every one litre of water.
- If Thrips, Redspidermite or Powdery mildew are already present add 15ml of product to every one litre of water.
- Spray thoroughly.
- Wash out sprayer after use.
- Do not store surplus spray mixture.