



STERISEAL

REG. NO. L 4231 / N-AR 0154
W 130196

Act No. 36 of 1947

A FUNGICIDAL PRUNING WOUND
PASTE. FOR USE ON ROSES,
GRAPEVINES, SHRUBS AND FRUIT
TREES AFTER PRUNING

'N SWAMDODER SNOEIWONDMIDDEL.
VIR GEBRUIK OP ROSE, WINGERDE,
STRUIKE EN VRUGTEBOME NA SNOEI

Active Ingredients / Aktiewe Bestanddele

"Busan 1009".....10 g/kg....."Busan 1009"

8-Hydroxyquinoline sulphate.....10 g/kg.....8-Hidroksikinoliensulfaat

NET VOLUME / NETTO VOLUME

REGISTRATION HOLDER / REGISTRASIEHOUER
AGRO-SERVE (Pty) Ltd trading as EFEKTO

REG. NO. / REG. NR.
1973/000868/07

Registered and Distributed by / Geregistreeer en Versprei deur:
Agro-Serve (Pty) Ld trading as EFEKTO™ Reg. No. 1973/000868/07 Private Bag X 215, Bryanston, 2021
EFEKTO and the EFEKTO logo are trademarks of Agro-Serve (Pty) Ltd.
© Agro-Serve (Pty) Ltd. All rights reserved.

CUSTOMER CARE / KLIËNTEDIENS: Tel: 0861 EFEKTO
customercare@efekto.co.za / www.efekto.co.za

BATCH NUMBER / LOTNOMMER

DATE MANUFACTURED / DATUM VERVAARDIG

UN NUMBER: 2902



SKADELIK / HARMFUL

WAARSKUWINGS

Bêre weg van voedsel en voer.

Hou buite bereik van kinders, oningeligte persone en diere.

VOORSORGMAATREËLS

Was met seep en koue water na gebruik.

Voorkom besoedeling van voedsel, eetgerei en drinkwater.

Vernietig leë houers en moet vir geen ander doel hergebruik nie.

GEBRUIKSAANWYSINGS

GEBRUIK SLEGS SOOS AANGEDUI

Skud deeglik en wend onmiddellik na snoei of afsaag aan op snoeivlakke en -wonde.

NOTA:

1. **STERISEAL** verminder besmetting van snoeiwonde en vlakke deur die volgende terugsterf swamme : *Eutypa armeniaca*, *Stereum purpureum*, *Polyporus hirsutus*, *P. versicolor* en *P. adustus*; en
2. Die terugsterf bakterieë *Pseudomonas syringae*. Dit verskaf 'n beperkte vermindering in die moontlikheid van besmetting deur *Xanthomonas pruni*.

WARNINGS

Store away from food and feed.
Keep out of reach of children, uninformed persons and animals.

PRECAUTIONS

Wash with soap and cold water after use.
Prevent contamination of food, feed, utensils and drinking water.
Destroy empty container and do not re-use for any other purpose.

DIRECTIONS FOR USE

USE ONLY AS DIRECTED

Shake well and apply to pruned surfaces and wounds immediately after pruning or cutting.

NOTE :

1. **STERISEAL** reduces infection of pruning wounds and cut surfaces by the following die-back fungi : *Eutypa armeniaca*, *Stereum purpureum*, *Polyporus hirsutus*, *P. versicolor* and *P. adustus*; and
2. The die-back bacteria *Pseudomonas syringae*, also provides a limited reduction in the possibility of infection by *Xanthomonas pruni*.